

The stromal vascular fraction from fat tissue in the treatment of osteochondral knee defect: Case report

Teplov O., Galimov D., Plakseichuk Y., Yagudin R., Pankov I., Rizvanov A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Salikhov, Masgutov, Chekunov, Tazetdinova, Masgutova, Teplov, Galimov, Plakseichuk, Yagudin, Pankov and Rizvanov. In this study we applied autologous fat tissue stromal vascular fraction (SVF) cells in combination with microfracturing technique in a 36-year-old man with an osteochondral lesion of the medial femoral condyle 8 months after the injury. Cell material was generated by fat tissue liposuction from the anterior abdominal wall with subsequent extraction of the SVF and injected through a mini-arthrotomy portal with subsequent fibrin sealant fixation. The follow-up period was 2 years. Clinical score improved from 23 to 96 according to IKDC and from 10 to 90 according to EQ-VAS at 24 months follow-up. Magnetic resonance imaging (MRI) before the surgery revealed an osteochondral lesion with development of significant trabecular edema that remained unchanged for 6 months despite conservative treatment. MRI 1 and 2 years after the surgery showed the recovery of the damaged cartilage thickness with somewhat uneven structure and a decrease in the trabecular edema of the femoral condyle. The use of SVF cells with fibrin sealant fixation might be a promising approach in the treatment of osteochondral joint lesions. Further studies are required.

<http://dx.doi.org/10.3389/fmed.2018.00154>

Keywords

Fat tissue, Fibrin sealant, Mesenchymal stem cells, Osteochondral lesions of the knee, Stromal vascular fraction

References

- [1] Chamberlain G, Fox J, Ashton B, Middleton J. Concise review: mesenchymal stem cells: their phenotype, differentiation capacity, immunological features, and potential for homing. *Stem Cells* (2007) 25:2739-49. doi: 10.1634/stemcells.2007-0197
- [2] Enea D, Cecconi S, Calcagno S, Busilacchi A, Manzotti S, Kaps C, et al. Single-stage cartilage repair in the knee with microfracture covered with a resorbable polymer-based matrix and autologous bone marrow concentrate. *Knee* (2013) 20:562-9. doi: 10.1016/j.knee.2013.04.003
- [3] Filardo G, Perdisa F, Roffi A, Marcacci M, Kon E. Stem cells in articular cartilage regeneration. *J Orthop Surg Res.* 11:42. doi: 10.1186/s13018-016-0378-x
- [4] Freitag J, Bates D, Boyd R, Shah K, Barnard A, Huguenin L, et al. Mesenchymal stem cell therapy in the treatment of osteoarthritis: reparative pathways, safety and efficacy-a review. *BMC Musculoskelet Disord.* (2016) 17:230. doi: 10.1186/s12891-016-1085-9
- [5] Yuk-wai Lee W, Wang B. Cartilage repair by mesenchymal stem cells: clinical trial update and perspectives. *J Orthopaed Trans.* (2017) 9:76-88. doi: 10.1016/j.jot.2017.03.005

- [6] Schinhan M, Gruber M, Vavken P, Dorotka R, Samouh L, Chiari C, et al. Critical-size defect induces unicompartmental osteoarthritis in a stable ovine knee. *J Orthop Res.* (2012) 30:214-20. doi: 10.1002/jor.21521
- [7] Moyad TF. Cartilage Injuries in the Adult Knee. *Eval Manag. Cart.* (2011) 2:226-36. doi: 10.1177/1947603510383973
- [8] Masgutov RF, Salihov RZ, Plaseichuk YA, Salafutdinov II, Rizvanov AA, Bogov AA. Use of stromal vascular fraction of adipose tissue for treatment of femur pseudoarthrosis: case report. *Cell Transplant Tiss Eng.* (2013) 8:116-8
- [9] Solovyeva VV, Salafutdinov II, Tazetdinova LG, Masgutov RF, Khaiboullina SF, Rizvanov AA. Genetic modification of adipose derived stem cells with recombinant plasmid DNA pBud-VEGF-FGF2 results in increased of IL-8 and MCP-1 secretion. *J Pure Appl Microbiol.* (2014) 8:523-528
- [10] Anderson AF, Irrgang JJ, Kocher MS, Mann BJ, Harrast JJ. The International Knee documentation committee subjective knee evaluation form: normative data. *Am J Sports Med.* (2006) 34:128-35. doi: 10.1177/0363546505280214
- [11] EuroQol Group. EQ-5D-5L User Guide. Rotterdam: EuroQol Group. (2011) Available online at: https://euroqol.org/wp-content/uploads/2016/09/EQ-5D-5L_UserGuide_2015.pdf (Accessed Apr 12, 2012)
- [12] Mithoefer K, McAdams T, Williams RJ, Kreuz PC, Mandelbaum BR. Clinical efficacy of the microfracture technique for articular cartilage repair in the knee: an evidence-based systematic analysis. *Am J Sports Med.* (2009) 37:2053. doi: 10.1177/0363546508328414
- [13] Gille J, Behrens P, Volpi P, de Girolamo L, Reiss E, Zoch W, et al. Outcome of Autologous Matrix Induced Chondrogenesis (AMIC) in cartilage knee surgery: data of the AMIC Registry. *Arch Orthop Trauma Surg.* (2013) 133:87-93. doi: 10.1007/s00402-012-1621-5
- [14] Behrens P. Matrixgekoppelte Mikrofrakturierung. *Arthroskopie.* (2005) 18:193-7. doi: 10.1007/s00142-00-0316-0
- [15] Verdonk P, Dhollander A, Almqvist KF, Verdonk R, Victor J. Treatment of osteochondral lesions in the knee using a cell-free scaffold. *Bone Joint J.* (2015) 97-B:318-23. doi: 10.1302/0301-620X.97B3.34555
- [16] Huang JI, Kazmi N, Durbhakula MM, Hering TM, Yoo JU, Johnstone B. Chondrogenic potential of progenitor cells derived from human bone marrow and adipose tissue: a patient-matched comparison. *J Orthop Res.* (2005) 23:1383-9. doi: 10.1016/j.orthres.2005.03.008.1100230621
- [17] Zhou Q, Li B, Zhao J, Pan W, Xu J, Chen S. IGF-I induces adipose derived mesenchymal cell chondrogenic differentiation in vitro and enhances chondrogenesis in vivo. *In Vitro Cell Dev Biol Anim.* (2016) 52:356-364. doi: 10.1007/s11626-015-9969-9
- [18] Riordan NH, Thomas E, Ichim W-P, Wang H, Solano F, Lara F, et al. Non-expanded adipose stromal vascular fraction cell therapy for multiple sclerosis. *J Transl Med.* (2009) 7:29. doi: 10.1186/1479-5876-7-29
- [19] Bornes TD, Adesida AB, Jomha NM. Mesenchymal stem cells in the treatment of traumatic articular cartilage defects: a comprehensive review. *Arthr Res Ther.* (2014) 16:432. doi: 10.1186/s13075-014-0432-1
- [20] Gibbs N, Diamond R, Sekyere EO, Thomas WD. Management of knee osteoarthritis by combined stromal vascular fraction cell therapy, platelet-rich plasma, and musculoskeletal exercises: a case series. *J Pain Res.* (2015) 8:799-806. doi: 10.2147/JPR.S92090
- [21] Pak J, Lee JH, Park KS, Park M, Kang L-W, Lee SH. Current use of autologous adipose tissue-derived stromal vascular fraction cells for orthopedic applications. *J Biomed Sci.* (2017) 24:9. doi: 10.1186/s12929-017-0318-z
- [22] Adachi N, Ochi M, Deie M, Ito Y. Transplant of mesenchymal stem cells and hydroxyapatite ceramics to treat severe osteochondral damage after septic arthritis of the knee. *J Rheumatol.* (2005) 32:1615-8
- [23] Buda R, Vannini F, Cavallo M, Grigolo B, Cenacchi A, Giannini S. Osteochondral lesions of the knee: a new one-step repair technique with bone-marrow-derived cells. *J Bone Joint Surg Am.* (2010) 92:2-11. doi: 10.2106/JBJS.J.00813
- [24] Nguyen PD, Tran TD, Nguyen HT, Vu HT, Le PT, Phan NL, et al. Comparative clinical observation of arthroscopic microfracture in the presence and absence of a stromal vascular fraction injection for Osteoarthritis. *Stem Cells Transl Med.* (2017) 6:187-95. doi: 10.5966/sctm.2016-0023
- [25] Trattnig S, Winalski CS, Marlovits S, Jurvelin JS, Welsch GH, Potter HG. Magnetic resonance imaging of cartilage repair: a review. *Cartilage* (2011) 2:5-26. doi: 10.1177/1947603509360209
- [26] Alparslan L, Winalski CS, Boutin RD, Minas T. Postoperative magnetic resonance imaging of articular cartilage repair. *Semin Musculoskelet Radiol.* (2001) 5:345-63. doi: 10.1055/s-2001-19044